

Probing magnetic correlations in heavy fermion $\text{Ce}M\text{In}_5$ ($M=\text{Rh, Ir, Co}$) with neutron scattering

W. Bao¹, P.G. Pagliuso¹, A.D. Christianson¹, J.L. Sarrao¹, J.D. Thompson¹, G. Aeppli², Z. Fisk⁶, J.S. Gardner⁵, J.W. Lynn³, S. Raymond⁴, A.H. Lacerda¹

¹ Los Alamos National Laboratory, Los Alamos, NM 87544, USA

² NEC Research, 1 Independent Way, Princeton, NJ 08540, USA

³ National Institute of Standard and Technology, Gaithersburg, MD 20899, USA

⁴ CEA-Grenoble, 38054 Grenoble Cedex 9, France

⁵ Chalk River Laboratories, Chalk River, Ontario K0J 1J0, Canada

⁶ Florida State University, Tallahassee, FL 32306, USA

Antiferromagnetism and superconductivity, with very high transition temperatures for heavy fermion compounds, exist in close proximity in the heavy fermion materials with chemical formula $\text{Ce}M\text{In}_5$ ($M=\text{Rh, Ir, Co}$). These tetragonal materials consist of alternating layers of the cubic heavy fermion antiferromagnet CeIn_3 and the intervening $M\text{In}_2$. Magnetic structures of the heavy fermion antiferromagnets CeRhIn_5 , Ce_2RhIn_8 , $\text{Ce}(\text{Rh, Ir})\text{In}_5$, and $(\text{Ce, La})\text{RhIn}_5$ are determined using neutron diffraction. Dynamic magnetic correlations of CeRhIn_5 , CeIrIn_5 and CeCoIn_5 are investigated with inelastic neutron scattering. Effects of hydrostatic pressure and magnetic field on the antiferromagnetic orders are also investigated. Insight on relation between the quasi-two-dimensional crystal structure, antiferromagnetism and superconductivity in these new heavy fermion materials, obtained through systematic study of the family with various M , changing ratio of the CeIn_3 and $M\text{In}_2$ layers, doping on the Ce site, and external fields, will be discussed.